



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

No. 2. Example: Given $x^2 - 19y^2 = 1$, to find x and y .

Value N m^2 $n^2 + \text{Diff.}$ $n^2 - \text{Diff.}$

$$19 = 19 \times 1^2 = 4^2 + 3 = 5^2 - 6$$

$$76 = 19 \times 2^2 = 8^2 + 12 = 9^2 - 5$$

$$171 = 19 \times 3^2 = 13^2 + 2$$

This gives at once $\left\{ \begin{array}{l} m=3 \\ n=13 \\ \text{Diff. } +2 \end{array} \right\}$ and in (A) we have $y=39$,
 $x=180$.

Proof: $28900 - 19 \times 1521 = 1$.

No. 3. Example: $x^2 - 31y^2 = 1$, to find x and y .

Value N m^2 $n^2 +$ Diff. $n^2 - \text{Diff.}$ Terms No. 1 2 3

$$31 = 31 \times 1 = 5^2 + 6^2 = 6 - 5$$

$$124 = 31 \times 2^2 = 11^2 + 3$$

$$279 = 31 \times 3^2 = \dots\dots\dots 17^2 - 10$$

$$496 = 31 \times 4^2 = 22^2 + 12$$

$$775 = 31 \times 5^2 = \dots\dots\dots 28^2 - 9$$

$$\left. \begin{array}{l} \text{Take } m = 3 + 2 + 5 \\ n = 17 + 11 + 28 \\ \text{Diff. } -10 + 3 - 2 \\ D_1 \quad +13 - 5 \\ D_2 \quad -18 \dots -18 \end{array} \right\}$$

$$\left. \begin{array}{l} 7 \\ 39 \\ -2 \end{array} \right\} \text{give } y=273 \\ x=1520$$

(TO BE CONTINUED.)

1. Proposed by EARL D. WEST, West Middleburg, Logan County, Ohio.

It is required to divide a given square number into two such parts that each part will be a square number.

I. Solution by Hon. JOSIAH H. DRUMMOND, LL. D., Portland, Maine.

No general solution can be given because there are comparatively few square numbers which are severally equal to the sum of two other squares. The well known equation $(2pq)^2 + (p^2 - q^2)^2 = (p^2 + q^2)^2$, in which $2pq$, $p^2 - q^2$, and $p^2 + q^2$ represent the sides of right triangles, is believed to include the cases in which the sum of two squares equals a third square. If this is true, it follows that the square root of the given square $(p^2 + q^2)^2$ must also be the sum of two squares, $p^2 + q^2$; and we have $p^2 + q^2 =$ the square root of the given number to find $(2pq)^2$ and $p^2 - q^2$, p being $> q$. If this can be done only by trial let us determine the limits within which the trials may be confined. Let m be the square root of the given number, and p and q may be any part positive numbers the sum of whose squares $= m$. Also, if any factor of m is equal to the sum of two squares, say s^2 and t^2 , p and q may be s and t multiplied, respectively, by the corresponding factor, but, if any value of p is not prime to the corresponding value of q , the value obtained by some of the processes may be the same as the value obtained by another process. If $p^2 + q^2 = m$, remembering that $p > q$, we have $p < \sqrt{m}$ and $p < \sqrt{\frac{m}{2}}$. Take $m=82$, then $p < 10$ and $p > 6$; that is, p may be 7, 8, or 9, but $=7$ or 8 gives impossible values for q ; while $p=9$, gives $q=1$. Then $2pq=18$ and $(p^2 - q^2)=80$. Hence 18^2 and 80^2 are the numbers, and we have $18^2 + 80^2 = 82^2$. But $82 = 2 \times 41$; then $s < 7$ and $s > 4$, and so may be 5

or 6; but $s=6$ makes t impossible, while $s=5$ makes $t=4$, and we have $2st=40$ and $(s^2-t^2)=18$. So $2pq=80$ and $(p^2-q^2)=18$, as before. Take $m=50$, p may be 6 or 7, but 6 makes q impossible, while $p=7$ gives $q=1$, and $2pq=14$ and $p-q^2=48$. $\therefore 14^2+48^2=50^2$. But $50=5 \times 10$, and $5=4+1$; then $s=2$ and $t=1$, $2st=4$, and $s^2-t^2=3$, $2pq=40$, and $p^2-q^2=30$, and $30^2+40^2=50^2$. Solutions may be obtained from other factors, but they give one of these two results. If $m=65$, it may be shown in the same manner that there are 4 pairs of numbers answering the conditions.

II. Solution by R. J. ADCOCK, Larchland, Illinois.

Since $(x^2+y^2+z^2+u^2+v^2)^2 = (x^2+y^2+z^2+u^2+v^2)^2 + (2xv)^2 + (2yv)^2 + (2zv)^2 + (2uv)^2$, is true for two or any greater number of letters. Then

$$\left[(2x+1)^2 + (2y)^2 \right]^2 = \left[(2x+1)^2 - (2y)^2 \right]^2 + \left[2(2x+1)2y \right]^2.$$

$$\text{And, } 1 = \left[\frac{(2x+1)^2 - (2y)^2}{(2x+1)^2 + (2y)^2} \right]^2 + \left[\frac{2(2x+1)2y}{(2x+1)^2 + (2y)^2} \right]^2.$$

If n^2 be the given square number,

$$n^2 = \left[\frac{(2x+1)^2 - (2y)^2}{(2x+1)^2 + (2y)^2} \times n \right]^2 + \left[\frac{2(2x+1)2yn}{(2x+1)^2 + (2y)^2} \right]^2 \text{ are the}$$

two square parts into which n^2 is divided. Observing that x may have any value including 0, y any value not = 0, and in order to avoid repetition of parts into which the square number is divided, the numbers for x and y must not make $2x+1$ and $2y$ have a common factor or be equal. If $x=0$, and $y=1$, the two square parts of n^2 are $\left(\frac{3n}{5}\right)^2$ and $\left(\frac{4n}{5}\right)^2$.

Also solved by H. W. Draughton, G. B. M. Zerr, P. S. Berg, A. L. Foote, P. H. Philbrick, and H. C. Whitaker.

2. Proposed by J. M. COLAW, Principal of High School, Monterey, Virginia.

Find two numbers, such that the difference of their squares may be a cube, and the difference of their cubes a square.

I. Solution by G. B. M. ZERR, A. M., Principal of High School, Staunton, Virginia.

Let $6x^3$, $10x^3$ be the numbers.

Then $100x^6 - 36x^6 = 64x^6 = \text{a cube number}$, and $1000x^9 - 216x^9 = 784x^9 = (28)^2 x^9 = \text{a square number}$, when x is a perfect square. Let $x=1$, then 6 and 10 are the numbers. Other values can be obtained by substituting 4, 9, 16, etc. for x .

II Solution by C. W. M. BLACK, Department of Mathematics in Wilmington Conference Academy, Dover, Delaware.

(A). Represent the numbers by x , and $x+a$. The difference of the cubes will be (1), $a(3x^2+3ax+a^2)$, and the difference of the squares (2), $a(2x+a)$. For one solution the first will be a square if $a=3x^2+3ax+a^2$, and the second a cube if $a=(a+2x)^2$. Solving these equations, $a=1$ and $x=-1$ or 0, $a=0$ and $x=0$, making the required numbers -1 and 0, 0 and 1, or 0 and 0. Also, $a(2x+a)$ will be a cube if (3), $a^2=a+2x$. Combining (1) and (3), $(a-1)(3a^2+3a+4)=0$. The first factor gives $a=1$, whence $x=-1$ or 0, as before. The second factor gives an imaginary result. As these results are not satisfactory, we seek another method for finding other values, if there are any.

(B). It may easily be shown that if x and $x+a$ are positive integers